



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

*Foreign and insular statistical reports of countries and cities—Yearly and monthly.*

AFRICA—*Lourenço Marquez*.—Month of September, 1902. Estimated population, 7,000. Total number of deaths, 30, including 6 from phthisis pulmonalis.

BAHAMAS—*Dunmore Town*.—Two weeks ended November 7, 1902. Estimated population, 1,232. Three deaths. No contagious diseases.

*Governors Harbor*.—Week ended November 8, 1902. Estimated population, 1,500. No deaths and no contagious diseases.

*Green Turtle Cay—Abaco*.—Two weeks ended October 29, 1902. Estimated population, 3,314. No deaths and no contagious diseases.

*Nassau*.—Two weeks ended October 8, 1902. Estimated population, 12,535. Number of deaths not reported. No contagious diseases reported.

BRAZIL—*Pernambuco*.—Two weeks ended September 30, 1902. Estimated population, 200,000. Total number of deaths, 194, including enteric fever, 4; whooping cough, 1; smallpox, 3, and 36 from phthisis pulmonalis.

CANADA—*British Columbia—Vancouver*.—Month of October, 1902. Estimated population, 29,000. Total number of deaths not reported. No deaths from contagious diseases reported.

*Ontario—Hamilton*.—Month of October, 1902. Estimated population, 54,000. Total number of deaths, 55, including diphtheria, 1; enteric fever, 1, and 5 from tuberculosis.

*Quebec—Sherbrooke*.—Month of October, 1902. Estimated population, 12,000. Total number of deaths, 15, including 1 from diphtheria.

GREAT BRITAIN—*England and Wales*.—The deaths registered in 76 great towns in England and Wales during the week ended October 18, 1902, correspond to an annual rate of 17.4 per 1,000 of the aggregate population, which is estimated at 14,862,163.

*Bradford*.—Two weeks ended October 18, 1902. Estimated population, 281,770. Total number of deaths, 184, including diphtheria, 2; enteric fever, 1; measles, 1; scarlet fever, 3; whooping cough, 1, and 16 from tuberculosis.

*London*.—One thousand four hundred and ninety-six deaths were registered during the week, including measles, 31; scarlet fever, 10; diphtheria, 21; whooping cough, 25; enteric fever, 24, and diarrhea, 60. The deaths from all causes correspond to an annual rate of 17.0 per 1,000. In Greater London 1,954 deaths were registered. In the "outer ring" the deaths included 5 from diphtheria, 6 from measles, 1 from scarlet fever, and 3 from whooping cough.

*Ireland*.—The average annual death rate represented by the deaths registered during the week ended October 18, 1902, in the 21 principal town districts of Ireland was 22.8 per 1,000 of the population, which is estimated at 1,092,401. The lowest rate was recorded in Armagh,

viz, 6.9, and the highest in Kilkenny, viz, 49.1 per 1,000. In Dublin and suburbs 200 deaths were registered, including diphtheria, 1; enteric fever, 2; scarlet fever, 2; whooping cough 1, and 26 from tuberculosis.

*Scotland*.—The deaths registered in 8 principal towns during the week ended October 18, 1902, correspond to an annual rate of 18.0 per 1,000 of the population, which is estimated at 1,679,923. The lowest mortality was recorded in Leith, viz, 8.6, and the highest in Paisley, viz, 30.1, per 1,000. The aggregate number of deaths registered from all causes was 501, including diphtheria, 8; measles, 2; scarlet fever, 8, and 11 from whooping cough.

*ITALY—Genoa*.—Month of September, 1902. Estimated population, 221,774. Total number of deaths, 334, including enteric fever, 5; measles, 3; scarlet fever, 1, and 51 from tuberculosis.

*JAMAICA—Kingston*.—Month of October, 1902. Estimated population, 46,542. Total number of deaths, 135, including 12 from phthisis pulmonalis.

*PHILIPPINE ISLANDS—Manila*.—Month of July, 1902. Population, 302,154. Total number of deaths, 2,223, including enteric fever, 22; whooping cough, 5; cholera, 1,066, and 107 from tuberculosis.

*RUSSIA—Riga*.—Month of August, 1902. Estimated population, 282,000. Total number of deaths, 521, including diphtheria, 3; enteric fever, 6; measles, 7; scarlet fever, 15; whooping cough, 3, and 48 from phthisis pulmonalis.

*WEST INDIES—St. Thomas*.—Two weeks ended October 24, 1902. Estimated population, 11,012. Total number of deaths, 7. No contagious diseases.